

Suhailiza S<sup>1</sup> Kannan TP<sup>2,3,\*</sup>,  
Sarliza Yasmin S<sup>2</sup>

## Genetic Basis of Supernumerary Tooth

<sup>1</sup>Department of Paediatric  
Dentistry, Hospital Raja  
Permaisuri Bainun, 30990  
Ipoh, Perak, Malaysia

<sup>2</sup>School of Dental Sciences,  
Universiti Sains Malaysia  
16150 Kubang Kerian,  
Kelantan, Malaysia

<sup>3</sup>Human Genome Centre,  
School of Medical Sciences,  
Universiti Sains Malaysia  
16150 Kubang Kerian,  
Kelantan, Malaysia

**Abstract** - Odontogenesis is a complex process regulated by both genetic and molecular controls. The development of a tooth in the embryo stage is controlled by a series of signals which occur between tooth-forming epithelium and neural crest-derived ectomesenchyme. Though many genes are involved in tooth formation involving major signalling molecules, the bone morphogenetic protein and fibroblast growth factor are the most important ones involved in odontogenesis. Supernumerary tooth occurs because of imbalance in the expression of the signalling pathways and their inhibitors. This review highlights the various signalling molecules that play a role in odontogenesis in order to provide a better understanding of the molecular mechanisms involved in the formation of supernumerary tooth in humans.

Received 30 Dec 2018.  
Revised 15 Jan 2019.  
Accepted 30 Jan 2019.  
Published Online 27 June 2019

**Keywords** - Genetics, odontogenesis, signalling molecules, supernumerary tooth

\*Corresponding author:  
Thirumulu Ponnuraj Kannan  
E-mail: [kannan@usm.my](mailto:kannan@usm.my)

### 1 INTRODUCTION

Teeth are highly mineralised tissues located at the entrance of the alimentary tract in both invertebrates and vertebrates [1]. Teeth are the elements of dermal skeleton that are present in a wide range of jawed vertebrates [2]. Though the main function of teeth is in chewing food, yet, they are also associated with defence, display of dominance as well as in the vocalisation in humans [3]. The human dentition comprises 20 teeth in the primary dentition and 32 in the permanent dentition [4, 5, 6]. Tooth agenesis denotes missing tooth/teeth as a result of developmental failure that results in decreased number of normal complement in human dentition [7]. Conversely, a supernumerary tooth denotes any tooth or odontogenic structure that is formed from a tooth germ resulting in more than the usual number of any given region in a dental arch [4, 5, 6].

A search was made in databases using the keywords 'supernumerary teeth, mouse, humans, genetics'. The articles collected were subjected to a systematic review to analyse the genetic basis of supernumerary teeth. Supernumerary teeth can be seen in many genetic disorders; but they are more common in syndromes like Gardner's syndrome, cleft lip and palate and cleidocranial dysplasia (CCD) and less commonly seen in

Fabry disease, Nance-Horan syndrome, Ellis-Van Creveld syndrome, Rubinstein-Taybi syndrome and trichorhinophalangeal syndrome [8]. Genetic entities that represent supernumerary teeth as a salient finding have been attributed to autosomal dominant inheritance, X chromosome inheritance and to both the inheritance patterns based on their locus heterogeneity [9]. Also, there are many reports supporting the theory of familial tendency to supernumerary teeth which were more evident in the relatives of the affected individuals [10]. Moreover, Seema Gupta and Praveen Kumar reported based on their study that in 8.6% of cases, there was a history of the same abnormality observed in other members of the family, which ascertained the hereditary nature of hyperdontia to occur [11].

### 2 CONTROLLING MECHANISM OF TOOTH DEVELOPMENT

Teeth develop due to progressive interactions between the ectoderm and the underlying ectomesenchyme tissue [12]. Subsequently, the ectomesenchyme forms the dental laminae giving rise to human deciduous teeth at six weeks *in utero* [13]. The permanent successors, on the other hand, develop as lingual extensions of these primary laminae which occurs between 20 weeks *in utero* until the age of five [14]. The early

stage of tooth development or odontogenesis at the embryological stage is regulated by a series of signals occurring between tooth-forming epithelium and the neural crest-derived ectomesenchyme [12]. Each tissue layer then instructs the other to differentiate into incisors, canines, premolars, and molars [15]. The reciprocal interactions between ectoderm and ectomesenchyme regulate the different phases involved in odontogenesis [16] as well as in regulating the tooth number [17]. The different phases are **Initiation phase** which determines the tooth region and numbers; **morphogenesis phase** which determines the tooth type, size, shape, dimension and cusp number, and the **cytodifferentiation phase** which determines the tooth structure such as enamel and dentine formation and mineralization. Thus, odontogenesis is a complex process under tight control of genetic and molecular events.

Research on the roles of signals and tissue interactions in cultured tissue explants and in mutant mice have shown inductive signalling and hierarchies in downstream transcription factors during odontogenesis. The development of tooth occurs through a chain of signalling interactions between the oral epithelium as well as neural crest-derived mesenchyme, genetically controlled by various signalling molecules and pathways [18]. More than 200 genes have been found to have active functions in developing tooth. Most of the expression patterns can be viewed in a comprehensive graphical database of gene expression profiles at <http://honeybee.helsinki.fi/toothexp> [19]. The roles of signalling molecules and the expression of homeobox genes in odontogenesis indicate a complementary interaction between the 'field' and 'clone' theories [15]. The 'field' theory was first proposed by Butler [20], who postulated that each tooth within a class, e.g. molars, develop number, shape, size, and order of development because it belongs to a common field [21]. Nevertheless, a field gradient exists depending on the position of tooth in the field. Clone theory was proposed by Osborne [22], who reported that a single pre-programmed cell clone is responsible for the development of a specific class of tooth [23].

Major signalling molecules involved in the regulation of tooth embryogenesis belong to the bone morphogenetic protein (BMP), fibroblast growth factor (FGF), sonic hedgehog (SHH), and wingless-type (WNT) families. BMPs and FGFs

are the most important molecules of odontogenesis that are expressed in both ectoderm and ectomesenchyme, whereas, SHHs and WNTs are expressed only in the ectoderm [24]. Supernumerary tooth and tooth agenesis occur due to imbalance in the expression of these four major signalling pathways and their inhibitors. Their roles in regulating odontogenesis, in turn, determine the tooth number and patterning [16]. Various genes that play a role in causing dental anomalies are presented in Table 1 and a diagrammatic representation of the genes and signalling pathway involved in odontogenesis is shown in Fig. 1.

Table 1: Genes involved in various dental anomalies

Gene	Mutation	Dental anomaly
<i>Msx1, Msx2, Dlx1, Dlx2</i>	Double mutant	Initiation stage arrest
<i>Lhx6, Lhx7, Gli1, Gli3</i>		
<i>Pax9, Lef1, Runx2, Barx1</i>	Null	Bud stage arrest
<i>Shh, Fgfr2b</i>		
<i>Activin βA</i>	Null	Bud stage arrest Lack of incisors and mandibular molars
<i>Ctip2</i>	Null	Late bell stage arrest
<i>Gli2</i>	Null	Abnormal maxillary incisors
<i>Gli3</i>	Heterozygous	Arrest maxillary incisor development
<i>Edar</i>	Downless	Small or absent enamel knot
<i>Fgf10</i>	Null	Small tooth germ
<i>Wnt/β catenin</i>	K14 conditional KO	Mishappen tooth bud
<i>Ectodin, Sostdc1/wise</i>	Null	Abnormal cusp
<i>Ectodin, Apc, Sp6, Lrp4, IFT88</i>	Null	Supernumerary teeth
<i>Gas1, Osr2, Sprouty2, Sprouty4</i>		
<i>Msx2, Lama3, Sp3, Sp6, Amelx</i>	Null	Enamel hypoplasia
<i>Enamelin, Mmp20</i>		
<i>Smoothened</i>	K14 conditional KO	
<i>Connexin 43</i>	Dominant negative	
<i>Periostin</i>	Null	Incisor enamel defect, thinner enamel
<i>Eda, Follistatin, Wnt3</i>	K14 transgenic	No enamel
<i>Ameloblastin, Gdnf</i>	Null	
<i>Noggin</i>	K14 transgenic	Abnormal ameloblast
<i>DSPP, Msx2</i>	Null	Dentinogenesis imperfecta
<i>DMP1, Sp3, Sp6</i>	Null	Abnormal dentin structure
<i>Msx2, Sp6</i>	Null	Root malformation
<i>Shh</i>	Ptc <sup>+/+</sup>	Shorter root
<i>Noggin</i>	K14 transgenic	Failed to form multiple root




Figure 1: Genes and signalling pathways involved in odontogenesis

### 3 MAJOR SIGNALLING MOLECULES INVOLVED IN THE FORMATION OF SUPERNUMERARY TOOTH

#### 3.1 Bone morphogenetic protein

One of the first signals identified in inductive interactions between epithelium and ectomesenchyme are growth factors belonging to the family of BMPs. The BMP family comprises a large group of proteins that are frequently expressed during tooth morphogenesis. For example, BMP2, BMP4 and BMP7 are expressed in the dental epithelium, BMP2 and BMP7 are expressed during the bud stage, and BMP4 is expressed during the thickening of the dental lamina. Also, BMPs can function as bidirectional signalling factors between the epithelium and ectomesenchyme. In ectomesenchyme, the BMPs stimulate the expression of transcription factors muscle segment homeobox 1 (MSX1), muscle segment homeobox 2 (MSX2), early growth response 1 (EGR1), and high mobility group (HMG) domain of the lymphoid enhancer-binding factor 1 (LEF-1) gene or transcription factor [21].

Among the BMP family, BMP4 is essential for normal tooth development. BMP4 is required to induce several target genes in the dental ectomesenchyme including *MSX1*. Any breakdown that occurs in these inductive interactions will arrest tooth development during the bud stage. The expression of BMP4 is initiated in the epithelium; however, the expression will then switch to the ectomesenchyme when inductive possibilities are

acquired from the latter which suggests the ability of this molecule to induce its own expression in ectomesenchymal cells. The intense mesenchymal expression of BMP4 during the bud stage can be linked to the subsequent transfer of the inductive ability to the epithelium which leads to the formation of enamel knot [21], and later, the supernumerary tooth. Ectodin is a BMP-antagonist which is widely expressed in developing tooth germ, but noticeably absent from enamel knots. Mice with lack of function of Ectodin displayed several anomalies including the presence of a supernumerary molar. Similarly, supernumerary tooth develops due to failure in normal *Ectodin*-mediated inhibition from the adjacent ectomesenchyme. Ectodin can inhibit WNT signalling and the correct modulation of this pathway is critical in determining the correct number of tooth formation [25]. Murashima-Suginami *et al.*, [26] reported that supernumerary teeth occurred due to increased BMP signalling in USAG-1-deficient mouse model.

#### 3.2 Fibroblast growth factor (FGF)

In mammals, the FGF family comprises of 19 growth factors (FGF 1–19). FGF plays a significant role in regulating the growth and morphogenesis of tooth germ. They regulate gene expression in ectomesenchyme and stimulate the epithelial cellular division and proliferation. These take place during the different stages of tooth development; the early phases of morphogenesis, early epithelial invagination which generates tooth bud, and during the

assessment of epithelial folds which generates the dental cuspids [27]. FGF4 and FGF9 in the primary enamel knot epithelium induce proliferation of both dental epithelium and ectomesenchyme, and then, later regulate the cuspid development, whereas FGF3 and FGF10 in the underlying ectomesenchyme stimulate only cell division in dental epithelium to form the dental papillae. These signals are required for the expression of *SHH* in the primary enamel knot epithelium [25]. Study on mice showed that intracellular FGF antagonists such as Sprouty (*SPRY*) genes are produced in response to FGF signalling and modulate the transduction in target cells [28]. *SPRY2* and *SPRY4* are expressed in the epithelium and ectomesenchyme of developing tooth respectively. Any functional breakdown of these genes results in the formation of extra tooth [24].

### 3.3 Sonic hedgehog (*SHH*)

Early signalling interactions between oral epithelium and neural crest cells are envisaged to establish information pattern along the developing dental axis [19]. At this stage, high *SHH* expression acts as a mitogen that is essential for normal proliferation of tooth bud because it invaginates into the underlying ectomesenchyme [19]. Hedgehog signal transduction through *SHH* may influence tooth number. In the absence of normal *SHH* signal transduction, tooth development will be arrested. Appropriate restriction of *SHH* activity is important to ensure the correct number of tooth formation in the right positions [29]. Once the early tooth bud is formed, continuous and reiterative signalling between epithelium and ectomesenchyme enables further growth and morphogenesis, with the bud stage progressing into the cap and bell stage [30]. Therefore, dysregulation of *SHH* activity plays a key role in the formation of supernumerary tooth [25].

*SHH* signals are mediated by the presence of primary cilia, that projects from the surface of all eukaryotic cells. Mutations in several genes including the ciliary protein IFT88/Polaris can result in changes in *SHH* signalling activity and the development of teeth in mice [25]. Ciliary protein IFT88/Polaris encodes essential functional components of the primary cilia. Besides, research [29], has shown that the upregulation of *SHH* activity in diastema mesenchyme can produce ectopic tooth in mutant mice which suggests that the *SHH* signalling may play a role in tooth position. Another link between

*SHH* signal transduction and the presence of additional teeth has been provided by runt-related transcription factor (*RUNX2*) in mutant mice. *RUNX2* is essential for the normal differentiation of bone-forming osteoblasts [31, 32]. In mice, *RUNX2* is expressed in mesenchymal compartment of a tooth and a complete loss of function is associated with arrested tooth development; however, in the heterozygous mutant, rudimentary supernumerary tooth formation takes place lingual to the first molar tooth germ. *RUNX2* transcription in the mesenchyme can repress *SHH* signalling in the epithelium. Thus, in the absence of adequate suppression of *SHH* transduction in these mice, additional teeth can develop in these regions [25]. Maisa Seppala and colleagues reported that *SHH* interacts at the molecular level with various other signalling pathways, Fgf and Wnt in particular, for normal progression of tooth development [33].

### 3.4 Wingless integrated (*WNT*)

*WNT* proteins form a large family of secreted ligands that activate several intracellular signalling pathways [34]. *WNT* signals drive multiple stages involved in odontogenesis, from the initiation stage until the tooth differentiation, which are broadly expressed in oral as well as dental epithelium. *WNT* pathways work through several mediators. For instance,  $\beta$ -catenin stabilization and activation of *LEF1* transcription factor activates the canonical signalling, crucial in normal tooth development [25]. *LEF1* is necessary for tooth development to progress beyond bud stage and inhibition of this *WNT* signalling pathway arrests odontogenesis. Evidence also suggests that the normal regulation of this pathway is important to determine the correct number of tooth formation [25]. Overexpression of *LEF1* in the oral epithelium in transgenic mice produced multiple invaginations in tooth forming regions [35].

There is a firm link associated between unregulated *WNT* signalling and hyperdontia in humans. In the case of Gardner's syndrome, an autosomal dominant disorder characterised by multiple adenomatous polyps of the colon and rectum, patients exhibited dental anomalies such as multiple supernumerary teeth, odontomas, and tooth impactions. The causative adenomatous polyposis coli (*APC*) tumour suppressor gene is a known inhibitor of *WNT* signalling [36]. The expression of ectodysplasin-A (*EDA*) gene is also regulated by *WNT* family of proteins [37]. If *WNT* signalling is blocked during the early bell stage

when secondary enamel knots form, the expression of *EDA* is reduced and the molars form with flattened cusps. Therefore, WNT signalling is important for the development of molar cusps [38], while, overexpression of *EDA* leads to the formation of supernumerary tooth. Thus, overexpression of canonical WNT signalling, through the loss of function of its inhibitors or by overexpression of its effectors, leads to the formation of supernumerary tooth [15]. Multiple teeth have also been seen in the molar field where  $\beta$ -catenin has been overexpressed in mice [36]. According to Bei [15], the number of teeth that can develop from the molar field appears to be restricted by WNT signalling. Supernumerary teeth and altered morphology of the molar crown have been reported in *WNT10A* null mice, though it results in tooth agenesis phenotype in humans [39].

#### 4 OTHER SIGNALLING MOLECULES

##### 4.1 Tumour necrosis factor pathway

Mutation in the gene encoding *EDA* ligand (e.g. *EDA1*) or *EDA*-receptor can result in disruption of *EDA* signalling [40], which then can lead to the formation of a component of the tumour necrosis factor (TNF) pathway. *EDA* signalling is active in organs that develop through signalling between epithelium and ectomesenchyme [25] [19]. The levels of *EDA* signalling are important to determine the tooth number. Overexpression of *EDA1* splice variant in the oral epithelium of transgenic mice produces supernumerary premolar-like tooth. Therefore, signals from epithelium are essential for the initiation of tooth development. Deficiency of *EDA* signalling results in hypodontia, while, too much *EDA* can produce supernumerary tooth [12].

##### 4.2 Runt-related transcription factor gene

Runt-related transcription factor (*RUNX2*) is a principal gene involved in bone and tooth development [41]. It is an osteoblast-specific transcription factor which is necessary for the differentiation of pluripotent mesenchymal cells into osteoblasts [42]. The presence of *RUNX2* in fully differentiated cells establishes the fact that *RUNX2* is also required in maintaining the full function of cells, especially those in bones [42]. *RUNX2* is also crucial in the formation of tooth. It is a key mesenchymal factor that influences tooth morphogenesis and the subsequent differentiation of ameloblasts, odontoblasts and osteoblasts lining the bone in the periodontal space [41, 43]. The length of *RUNX2* is 220 kb

and has eight exons belonging to the runt domain (*RUNX*) family of genes. The genes, namely *RUNX1*, -2 and -3, exhibit high amino acid homology. Their protein products form a heterodimer with the core-binding factor  $\beta$  (CBF- $\beta$ ). CBF- $\beta$  is required for the function of *RUNX2* in skeletal development, which allosterically enhances DNA binding by RUNX proteins at runt homology domain (RHD). Moreover, it plays an important role in stabilizing the RUNX proteins against proteolytic degradation by the ubiquitin-proteasome system [43]. Several studies [31, 44, 45, 46, 47, 48, 49], on mutational analysis of *RUNX2* in cleidocranial dysplasia (CCD) patients, have shown that mutations in *RUNX2* gene are accountable for this syndrome [31]. CCD is a syndrome that affects the development of bone and tooth. The most common features of CCD include delayed closure of skull sutures, hypoplastic or aplastic clavicles and multiple supernumerary teeth [50]. *RUNX2* gene dysfunction in tooth-forming cells may directly result in dental anomalies in CCD patients [43]. Therefore, these dental abnormalities suggest the important role that *RUNX2* plays during odontogenesis [51]. Under normal condition, *RUNX2* acts as a cell growth inhibitor in immature osteoblasts by supporting an exit from the cell cycle and promoting increased expression of osteoblast phenotype [52]. Hence, *RUNX2* regulates cell proliferation and may have a specific control of the dental lamina and the subsequent formation of successive dentitions [43]. *In-vitro* genetic studies have proved that deletion or deficiency of *RUNX2* in knockout mice arrested the tooth development at the bud or early cap stage, and the osteoblasts lining bone in the periodontal space [41, 53]. In contrast, another study has reported that loss of function of *RUNX2* gene would support the proliferation of dental lamina. For example, the reduced function of *RUNX2* gene caused the development of supernumerary tooth in CCD patients [42].

Studies showed that both *RUNX2*<sup>-/-</sup> and *RUNX2*<sup>+/-</sup> mice displayed lingual buds in front of upper molars, and these were much more prominent than those in the wild-type mice [54]. It was assumed that these buds represent the secondary dentition and that *RUNX2* plays a role in inhibiting the formation of these buds [55]. It may appear contradictory that inhibition of *RUNX2* gene function may arrest primary tooth development but stimulates the formation of secondary teeth [56]. Nevertheless, it is normal for the same gene to have different effects at

different developmental stages during the process of embryogenesis [54]. Thesleff [41], proposed that humans possess the potential to develop a third dentition which is normally inhibited by RUNX2 gene. This has been confirmed by Wang *et al.*, [55] who showed that RUNX2 gene inhibits serial tooth formation [41, 57]. Analysing the regulations, expressions and functions of RUNX2 gene, particularly in non-syndromic patients with supernumerary tooth would enhance the understanding of tooth development in humans [41]. The lack of teeth, as well as formation of supernumerary teeth, were attributed to the mutations in *AXIN2* and *RUNX2*, respectively, which occurred due to Wnt/ $\beta$ -catenin signalling modulation in dental mesenchyme [58]. They also reported that increased mesenchymal Wnt/ $\beta$ -catenin signalling can result in the inhibition of tooth initiation.

#### 5 RELEVANCE OF MOUSE MODEL TO HUMANS

Jussila and Thesleff [59], based on the phenotypes of two syndromes, namely CCD and craniosynostosis syndrome, suggested that the potential for continued tooth replacement may be unlocked in humans. The presence of supernumerary teeth has been suggested to denote a third dentition in CCD and in craniosynostosis syndrome due to mutations in the transcription factor *RUNX2* and the interleukin receptor *IL11RA*, respectively [60, 61]. However, in the mouse models of these syndromes, there are no supernumerary teeth; this could be attributed to the reason that the teeth in mouse are not normally replaced, and hence, unsuitable for studies involving tooth replacement [61, 62]. In mice, the number of teeth is lesser compared to humans and moreover, since mice have only a primary dentition, mouse models may not reflect the same to determine the cause of supernumerary teeth in humans [63]. Though in mouse models as well as in human syndromes, supernumerary teeth are induced by modulating signal pathways, this may not function in adult jaws, the reason being that these teeth are formed from the tissue associated with developing teeth which would not be present in the jaws of the adults [59]. D'Souza and Klein [64] reported that the use of multifaceted approaches involving mouse and human genetic researches were needed in order to reveal the precise aetiology of development of supernumerary tooth. Xi Lu and colleagues in their review reported that though the mouse dentitions were quite different

from that of humans, exploration of the molecular mechanism in mouse was still useful. However, they suggested that animals such as chimpanzee, which have more similarities in the development patterns with humans, need to be investigated to identify the genetic basis of supernumerary teeth [65].

#### 6 CONCLUSIONS

Recent studies have probed into the molecular mechanisms underlying tooth morphogenesis and differentiation. Although genetics may be implicated in the formation of supernumerary tooth, little is known about the initiation of tooth formation, the genetic regulation of successional teeth, as well as the underlying mechanisms involved in its formation. Nonetheless, a better understanding of the roles of these aforementioned signalling molecules, particularly WNT and RUNX2, will provide fundamental insights into the molecular genetics of supernumerary tooth in humans. This, in turn, may assist in future tooth regeneration and tooth engineering.

#### CONFLICTS OF INTEREST

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the paper.

#### ACKNOWLEDGEMENT

This review was supported by Universiti Sains Malaysia Short Term Grant 304/PPSG/61311087.

#### REFERENCES

- [1] Koussoulakou DS, Margaritis LH, Koussoulakos SL. A curriculum vitae of teeth: evolution, generation, regeneration. *International Journal of Biological Sciences*. 2009;5(3):226-243.
- [2] Huysseune A, Sire JY, Witten PE. Evolutionary and developmental origins of the vertebrate dentition. *Journal of Anatomy*. 2009;214(4):465-476.
- [3] Chinsebu KC. Teeth are bones: signature genes and molecules that underwrite odontogenesis. *JMGG*. 2012;4(2):13-24.
- [4] Schulze C. Developmental abnormalities of the teeth and jaws. 1970. in: Gorlin, R.J., Goldman, H.M. (Eds.), *Thomas's Oral Pathology*. Mosby Company, St Louis, pp. 96-183.
- [5] Scheiner MA, Sampson WJ. Supernumerary teeth: a review of the literature and four case reports. *Australian Dental Journal*. 1997;42(3):160-165.
- [6] Rajab LD, Hamdan MA. Supernumerary teeth: review of the literature and a survey of 152 cases. *International Journal of Paediatric Dentistry*. 2002;12(4):244-254.
- [7] Pani SC. The genetic basis of tooth agenesis: basic concepts and genes involved. *Journal of the Indian*

- Society of Pedodontics and Preventive Dentistry. 2011;29(2):84-89.
- [8] Subasioglu A, Savas S, Kucukyilmaz E, Kesim S, Yagci A, Dundar M. Genetic background of supernumerary teeth. *European Journal of Dentistry*. 2015;9(1):153-158.
- [9] Cammarata-Scalisi F, Avendaño A, Callea M. Main genetic entities associated with supernumerary teeth. *Archivos Argentinos de Pediatría*. 2018;116(6):437-444.
- [10] Mallineni SK. Supernumerary Teeth: Review of the Literature with Recent Updates. *Conference Papers in Science*. 2014;2014: Article ID 764050, 6 pages.
- [11] Gupta S, Praveen Kumar PS. A study on prevalence, complications, and possible etiologic factors of supernumerary teeth in 6–12-year-old schoolchildren of Rohtak, India. *Indian Journal of Dental Sciences*. 2017;9:141-147.
- [12] Tucker AS, Sharpe P. The cutting edge of mammalian development; how the embryo makes teeth. *Nature Reviews Genetics*. 2004;5(7):499-508.
- [13] Sartaj R, Sharpe P. Biological tooth replacement. *Journal of Anatomy*. 2006;209(4): 503-509.
- [14] Ferguson CA, Tucker AS, Sharpe PT. Temporospatial cell interactions regulating mandibular and maxillary arch patterning. *Development*. 2000;127(2):403-412.
- [15] Bei M. Molecular genetics of tooth development. *Current Opinion in Genetics & Development*. 2009;19(5):504-510.
- [16] Brook AH. Multilevel complex interactions between genetic, epigenetic and environmental factors in the aetiology of anomalies of dental development. *Archives of Oral Biology*. 2009;54(Suppl 1): S3-S17.
- [17] Nakamura T, Fukumoto S. Genetics of supernumerary tooth formation. *Journal of Oral Biosciences*. 2013;55(4):180-183.
- [18] Williams MA, Letra A. The Changing Landscape in the Genetic Etiology of Human Tooth Agenesis. *Genes*. 2018;May16;9(5):pii: E255.
- [19] Cobourne MT, Sharpe PT. Tooth and jaw: molecular mechanisms of patterning in the first branchial arch. *Archives of Oral Biology*. 2003;48(1):1-14.
- [20] Butler PM. Studies of the mammalian dentition. *Proceedings of the Zoological Society of London*. 1939;109:1-36.
- [21] Galluccio G, Castellano M, La Monaca C. Genetic basis of non-syndromic anomalies of human tooth number. *Archives of Oral Biology*. 2012;57(7):918-930.
- [22] Osborn JW. Morphogenetic gradients: fields versus clones. In: Butler PM, Joysey KA. editors. *Development, function and evolution of teeth*. Academic Press; London: 1978. pp. 171–201.
- [23] Townsend G, Harries EF, Lesot H, Clauss F, Brook AH. Morphogenetic fields within the human dentition: a new clinically relevant synthesis of an old concept. *Archives of Oral Biology*. 2009;54s: S34-S44.
- [24] Klein OD, Minowada G, Peterkova R, Kangas A, Yu BD, Lesot H, Peterka M, Jernvall J, Martin GR. Sprouty genes control diastema tooth development via bidirectional antagonism of epithelial-mesenchymal FGF signalling. *Development Cell*. 2006;11(2):181-190.
- [25] Fleming PS, Xavier GM, DiBiase AT, Cobourne MT. Revisiting the supernumerary: the epidemiological and molecular basis of extra teeth. *British Dental Journal*. 2010;208(1):25-30.
- [26] Murashima-Suginami A, Takahashi K, Sakata T, Tsukamoto H, Sugai M, Yanagita M, Shimizu A, Sakurai T, Slavkin HC, Bessho K. Enhanced BMP signaling results in supernumerary tooth formation in USAG-1 deficient mouse. *Biochemical and Biophysical Research Communications*. 2008;369:1012-1016.
- [27] Thesleff I, Sharpe P. Signalling networks regulating dental development. *Mechanisms of Development*. 1997;67(2):111-123.
- [28] Mason JM, Morrison DJ, Basson MA. Sprouty proteins: multifaceted negative-feedback regulators of receptor tyrosine kinase signalling. *Trends in Cell Biology*. 2006;16(1):45-54.
- [29] Cobourne MT, Miletich I, Sharpe PT. Restriction of sonic hedgehog signalling during early tooth development. *Development*. 2004;131(12):2875-2885.
- [30] Jernvall J, Thesleff I. Reiterative signalling and patterning during mammalian tooth morphogenesis. *Mechanisms of Development*. 2000;92(1):19-29.
- [31] Mundlos S, Otto F, Mundlos C, Mulliken JB, Aylsworth AS, Albright S, Lindhout D, Cole WG, Henn W, Knoll JH, Owen MJ, Mertelsmann R, Zabel BU, Olsen BR. Mutations involving the transcription factor CBFA1 cause cleidocranial dysplasia. *Cell*. 1997;89(5):773-779.
- [32] Schwabe GC, Opitz C, Tinschert S, Mundlos S, Sharpe PT. Molecular mechanisms of tooth development and malformations. *Oral Biosciences & Medicine*. 2004;1(2):77-91.
- [33] Seppala M, Fraser GJ, Birjandi AA, Xavier GM, Cobourne MT. Sonic Hedgehog Signaling and Development of the Dentition. *Journal of Developmental Biology*. 2017;5(2):6.
- [34] Logan CY, Nusse R. The Wnt signalling pathway in development and disease. *Annual Review of Cell and Developmental Biology*. 2004;20:781-810.
- [35] Nakamura T, de Vega S, Fukumoto S, Jimenez L, Unda F, Yamada Y. Transcription factor epiprofin is essential for tooth morphogenesis by regulating epithelial cell fate and tooth number. *Journal of Biological Chemistry*. 2008;283(8):4825-4833.
- [36] Järvinen E, Salazar-Ciudad I, Birchmeier W, Taketo MM, Jernvall J, Thesleff I. Continuous tooth generation in mouse is induced by activated epithelial Wnt/beta-catenin signalling. *Proceedings of the National Academy of Sciences of the United States of America*. 2006;103(49):18627-18632.
- [37] Laurikkala J, Mikkola M, Mustonen T, Åberg T, Koppinen P, Pispä J, Nieminen P, Galceran J, Grosschedl R, Thesleff I. TNF signalling via the ligand-receptor pair ectodysplasin and edar controls the function of epithelial signalling centers and is regulated by Wnt and activin during tooth organogenesis. *Developmental Biology*. 2001;229(2):443-455.
- [38] Liu F, Chu EY, Watt B, Zhang Y, Gallant NM, Andl T, Yang SH, Lu MM, Piccolo S, Schmidt-Ullrich R, Taketo MM, Morrisey EE, Atit R, Dlugosz AA, Millar SE. Wnt/beta-catenin signalling directs multiple stages of tooth morphogenesis. *Developmental Biology*. 2008;313(1):210-224.
- [39] Yang J, Wang SK, Choi M, Reid BM, Hu Y, Lee YL, Herzog CR, Kim-Berman H, Lee M, Benke PJ, Lloyd KC, Simmer JP, Hu JC. Taurodontism, variations in tooth number, and misshapened crowns in Wnt10a null mice and human kindreds. *Molecular Genetics & Genomic Medicine*. 2015;3:40-58.

- [40] Sedano HO, Gorlin R. Familial occurrence of mesiodens. *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*. 1969;27(3):360-362.
- [41] Thesleff I. Developmental biology and building a tooth. *Quintessence International*. 2003;34(8):613-620.
- [42] Ducy P, Zhang R, Geoffroy V, Ridall A, Karsenty G. *Osf2/Cbfa1*: a transcriptional activator of osteoblast differentiation. *Cell*. 1997;89:747-754.
- [43] Camilleri S, McDonald F. *RUNX2* and dental development. *European Journal of Oral Sciences*. 2006;114(5):361-373.
- [44] Quack I, Vonderstrass B, Stock M, Aylsworth AS, Becker A, Brueton L, Lee PJ, Majewski F, Mulliken JB, Suri M, Zenker M, Mundlos S, Otto F. Mutation analysis of core binding factor A1 in patients with cleidocranial dysplasia. *American Journal of Human Genetics*. 1999;65(5):1268-1278.
- [45] Otto F, Kanegane H, Mundlos S. Mutations in the *RUNX2* gene in patients with cleidocranial dysplasia. *Human Mutation*. 2002;19(3):209-216.
- [46] Currall V, Clancy R, Dimond D, Amirfeyz R, Kershaw C, Gargan M. Cleidocranial dysplasia. *Current Orthopaedics*. 2007;21(2):159-162.
- [47] Kolokitha OE, Papadopoulou AK. Impaction and apical root angulation of the maxillary central incisors due to supernumerary teeth: combined surgical and orthodontic treatment. *American Journal of Orthodontics and Dentofacial Orthopedics*. 2008;134(1):153-60.
- [48] Li Y, Pan W, Xu W, He N, Chen X, Liu H, Darryl Quarles L, Zhou H, Xiao Z. *RUNX2* mutations in Chinese patients with cleidocranial dysplasia. *Mutagenesis*. 2009;24(5):425-431.
- [49] Ryoo HM, Kang HY, Lee SK, Lee KE, Kim JW. *RUNX2* mutations in cleidocranial dysplasia patients. *Oral Diseases*. 2010;16(1):55-60.
- [50] Tang S, Xu Q, Xu X, Du J, Yang X, Jiang Y, Wang X, Speck N, Huang T. A novel *RUNX2* missense mutation predicted to disrupt DNA binding causes cleidocranial dysplasia in a large Chinese family with hyperplastic nails. *BMC Medical Genetics*. 2007;8:82-88.
- [51] Jensen BL, Kreiborg S. Development of the dentition in cleidocranial dysplasia. *Journal of Oral Pathology & Medicine*. 1990;19(2):89-93.
- [52] Pratap J, Galindo M, Zaidi SK, Vradii D, Bhat BM, Robinson JA, Choi JY, Komori T, Stein JL, Lian JB, Stein GS, van Wijnen AJ. Cell growth regulatory role of *Runx2* during proliferative expansion of preosteoblasts. *Cancer Research*. 2003;63(17):5357-5362.
- [53] D'Souza RN, Åberg T, Gaikwad J, Cavender A, Owen M, Karsenty G, Thesleff I. *Cbfa 1* is required for epithelial-mesenchymal interactions regulating tooth development in mice. *Development*. 1999;126(13):2911-2920.
- [54] Wang XP, Åberg T, James MJ, Levanon D, Groner Y, Thesleff I. *RUNX2* (*CBFA1*) inhibits *Shh* signalling in the lower but not upper molars of mouse embryos and prevents the budding of putative successional teeth. *Journal of Dental Research*. 2005;84(2):138-143.
- [55] Wang XP, O'Connell DJ, Lund JJ, Saadi I, Kuraguchi M, Turbe-Doan A, Cavallesco R, Kim H, Park PJ, Harada H, Kucherlapati R, Maas RL. *Apc* inhibition of *Wnt* signalling regulates supernumerary tooth formation during embryogenesis and throughout adulthood. *Development*. 2009;136(11):1939-1949.
- [56] Wang XP, Fan J. Molecular genetics of supernumerary tooth formation. *Genesis*. 2011;49(4):261-277.
- [57] Jonason JH, Xiao G, Zhang M, Xing L, Chen D. Post-translational regulation of *RUNX2* in bone and cartilage. *Journal of Dental Research*. 2009;88(8):693-703.
- [58] Järvinen E, Shimomura-Kuroki J, Balic A, Jussila M, Thesleff I. Mesenchymal *wnt/beta-catenin* signaling limits tooth number. *Development*. 2018;145(4):dev158048.
- [59] Jussila M, Thesleff I. Signaling networks regulating tooth organogenesis and regeneration, and the specification of dental mesenchymal and epithelial cell lineages. *Cold Spring Harbor Perspectives in Biology*. 2012;4:a008425.
- [60] Jensen BL, Kreiborg S. Development of the dentition in cleidocranial dysplasia. *Journal of Oral Pathology & Medicine*. 1990;19:89-93.
- [61] Nieminen P, Morgan NV, Fenwick AL, Parmanen S, Veistinen V, Mikkola ML, Giraud A, Judd L, Arte S, Brueton LA, Wall SA, Mathijssen IM, Maher ER, Wilkie AO, Kreiborg S, Thesleff I. Inactivation of *IL11* signaling causes craniosynostosis, delayed tooth eruption and supernumerary teeth. *American Journal of Human Genetics*. 2011;89:67-81.
- [62] D'Souza RN, Åberg T, Gaikwad J, Cavender A, Owen M, Karsenty G, Thesleff I. *Cbfa1* is required for epithelial-mesenchymal interactions regulating tooth development in mice. *Development*. 1999;126:2911-2920.
- [63] Takahashi M, Hosomichi K, Yamaguchi T, Yano K, Funatsu T, Adel M, Haga S, Maki K, Tajima A. Whole-exome sequencing analysis of supernumerary teeth occurrence in Japanese individuals. *Human Genome Variation*. 2017;4:16046.
- [64] D'Souza RN, Klein OD. Unraveling the molecular mechanisms that lead to supernumerary teeth in mice and men: current concepts and novel approaches. *Cells Tissues Organs*. 2007;186(1):60-69.
- [65] Lu X, Yu F, Liu J, Cai W, Zhao Y, Zhao S, Liu S. The epidemiology of supernumerary teeth and the associated molecular mechanism. *Organogenesis*. 2017;13:3,71-82.